

FIRST AID

IF IN EYES:

• Hold eye open and rinse slowly and gently with water for 15-20 minutes.  
• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.  
• Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

• Take off contaminated clothing.  
• Rinse skin immediately with plenty of water for 15-20 minutes.  
• Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

• Call a poison control center or physician immediately for treatment advice.  
• Have person sip a glass of water if able to swallow.  
• Do not induce vomiting unless told to do so by the poison control center or physician.  
• Do not give anything by mouth to an unconscious person.

IF INHALED:

• Move person to fresh air.  
• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.  
• Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if inhaled. Causes moderate eye irritation. Avoid breathing dust. Remove and wash contaminated clothing before reuse. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves such as natural rubber. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with caution when applying to turf areas adjacent to any body of water. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr provides season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in established lawns and ornamental turf, including golf course fairways, roughs and tee boxes, and in established landscape ornamentals. This product also provides postemergent crabgrass control up to the 4-leaf stage.

SPREADER SETTINGS			
to apply at the listed pounds per 1,000 square feet			
FILL & SET SPREADER	1,66 per 1,000 Sq Ft	3,33 per 1,000 Sq Ft	6,67 per 1,000 Sq Ft
Ortho Broadcast	3	5	8
Earthway Broadcast	8	15	19
Precision Broadcast	3½	4½	6½
Scotts Broadcast/Speedy Green	3	4	6

FOR USE ON TURF

Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr provides season-long control of crabgrass and control or suppression of other annual grasses and broadleaf weeds in established lawns and ornamental turf, including golf course fairways, roughs and tee boxes. Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr will not harm nearby established ornamentals identified on this label as "Tolerant Ornamentals" when used according to label directions.

Apply this product only to seeded, sodded, or sprigged turf that is well established. Before the use of Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr, roots must be developed and the turf mowed at least twice to a uniform height. Injury may result if the turf is not well established, or stressed by weather, pests or other conditions.

Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr may be applied to established turfgrass species listed on this label. When applied as directed, the following turfgrass species are tolerant to this product:

Sure Green™ Grass & Weed Barrier™ is a trademark of Spra-Green Lawn Fertilizer, LLC (CS-03082010N) 0316(01)

COOL-SEASON GRASSES	USE PRECAUTIONS
Creeping Bentgrass <sup>1</sup> ( <i>Agrostis palustris</i> ) , Kentucky Bluegrass ( <i>Poa pratensis</i> ) , Fine Fescue <sup>2</sup> ( <i>Festuca rubra</i> ) , Tall Fescue ( <i>Festuca arundinacea</i> ) , Perennial Ryegrass ( <i>Lolium perenne</i> )	<sup>1</sup> Turfgrass injury may result from the use of this product on certain varieties of creeping bentgrass such as Cohansey, Carmen, Seaside and Washington. Not all varieties of creeping bentgrass have been tested. Do not apply this product to Colonial Bentgrass ( <i>Agrostis tenuis</i> ) varieties.  <sup>2</sup> Turfgrass injury may result from use of this product on the following varieties of Fescue such as Atlanta, Banner, Beauty, Bilgart, CF-2, Enjoy, HF-93, Highlight, Inalo, Koket, Jamestown, Majenta, Mary, Pennlawn, Tamara, Tatiana, Walfort and Waldina. Not all varieties of fescue have been tested.
WARM-SEASON GRASSES	
Bahiagrass ( <i>Paspalum notatum</i> ) , Bermudagrass ( <i>Cynodon dactylon</i> ) , Buffalograss <sup>3</sup> ( <i>Buchloe dactyloides</i> ) , Carpetgrass ( <i>Axonopus compressus</i> ) , Centipedegrass ( <i>Eremochloa ophiuroides</i> ) , Kikuyugrass ( <i>Pennisetum clandestinum</i> ) , St. Augustinegrass ( <i>Stenotaphrum secundatum</i> ) , Zoysiagrass ( <i>Zoysia japonica</i> )	<sup>3</sup> Do not use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response to turfgrass varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

RESEEDING, OVERSEEDING OR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 12 weeks after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, use mechanical or power seeding equipment (silt seeders) for good seed-to-soil contact.

- USE RESTRICTIONS
- Do not apply more than 6.67 lbs./1,000 sq. ft. (0.5 lbs. AI/Acre) per application.
  - Do not apply more than 20 lbs./1,000 sq. ft. (1.5 lbs. AI/Acre) per year.
  - Do not use clippings from treated turf for animal feed or for mulching around vegetables or fruit trees
  - Do not allow domestic animals to feed or forage on treated turf.
  - Keep people and pets off treated areas until dust has settled.
  - Do not use on golf course putting greens.

CRABGRASS CONTROL: Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of crabgrass from the ground. Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr provides "early post-emergence" control during the early stages of growth after crabgrass emerges from the ground. This early post-emergence control is limited and applications are only effective if applied before the crabgrass has its 3-leaf or first tier (which usually is about the time when you can first see the crabgrass in the lawn). With this post-emergence activity, the applicator has 2 to 4 weeks longer to make applications for control of crabgrass.

APPLICATION INSTRUCTIONS: For best results, apply Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr within a few days after mowing and delay mowing again for a few days after application. Rainfall or irrigation activates the product. Improved weed control will result if the application is followed by ½ inch of rainfall or sprinkler irrigation. Weed control may be hampered if rainfall or irrigation does not occur with 30 days after application. Split, or half rate applications spaced 5 to 10 weeks apart may also enhance weed control. Cultural practices such as aeration and de-thatching should be completed and the turf allowed to recover before application of this product.

APPLICATION TIMING: This product may be applied as a single application, split application, or as a sequential application for crabgrass control in the spring, summer, or fall as described below.

SPRING APPLICATIONS: For single applications in the spring or early summer, follow the appropriate rate listed in the "Application Rates" table below. Rates are based on the location to be used (Region), the mowing height of the turfgrass, and whether the timing of the application is considered preemergence or postemergence. The length of residual weed control is dependent on the total rate applied, but may vary depending on weather, weed pressure and location.

Program 1: For preemergence control to turf mowed relatively high, such as home lawns, this program will provide 3 to 5 months of preemergence crabgrass control. Do not use this program for early postemergence control of crabgrass.

Program 2: For preemergence control at sites where the turf is mowed relatively low, such as golf course fairways, and where turfgrass maintenance and weed control applications were not made the previous year, this program will provide 4 to 6 months of preemergence crabgrass control and may also be used for early postemergence control of crabgrass at locations where turf is cut high.

Program 3: For preemergence control at locations where turf is mowed relatively low, such as golf course fairways, and where turfgrass maintenance and weed control applications were not made the previous year, this program will provide 4 to 6 months of preemergence crabgrass control and may also be used for early postemergence control of crabgrass at locations where the turf is cut relatively low.

FALL APPLICATIONS: Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr may be applied in late summer or early fall (late August through November). Follow the instructions for "Program 3" for crabgrass control through the early part of the next spring. Fall applications may be followed by an appropriate spring application to provide season-long control.



# GRASS & WEED BARRIER™

## Plus LAWN FOOD

### Containing Dithiopyr

For Season-Long Control of Crabgrass and Control or Suppression of Listed Annual Grasses and Broadleaf Weeds in Established Lawns and Ornamental Turf, including Golf Courses, Fairways, Roughs and Tee Boxes and in Established Landscape Ornamentals.

- Controls and Suppresses over 16 Grasses and 23 Broadleaf Weeds
- Provides Pre and Post Emergence Control of Crabgrass
- Apply Spring and Fall for Optimal Crabgrass Results
- Covers Up to 12,000 Sq. Ft. in Southern Lawns
- For Use on Listed Landscape Ornamental Plants

ACTIVE INGREDIENT: Dithiopyr: S,S'-dimethyl 2-(di-fluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarboximate.....0.172% OTHER INGREDIENTS.....99.828% TOTAL.....100.00%

EPA REG. NO. 53883-209-87346 EPA EST. NO. 87346-NM-001 Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for further precautions and First Aid information.

Distributed by:Guaranteed by:  
Spra-Green Lawn Fertilizer, LLC  
PO BOX 845 • 221 N. Main • Portales, NM 88130  
Sure Green™ Grass & Weed Barrier™ is a trademark of Spra-Green Lawn Fertilizer, LLC (CS-03082010N) 0316(01)

NET WT. 40 LBS. (18.14 KG)

Turf Grass Regions			
Table 1: Application Rates For Crabgrass Control			
REGION	PROGRAM 1 Preemergence (high-cut turf)	PROGRAM 2 Preemergence (low-cut turf) Postemergence (high-cut turf)	PROGRAM 3 Preemergence (low-cut turf) Postemergence (low-cut turf)
North(all areas)	1.66 lbs./1,000 sq. ft. (0.125 lbs. AI/acre)	2.45 lbs./1,000 sq. ft. (107 lbs./acre or 0.183 lbs. AI/acre)	3.33 lbs./1,000 sq. ft. (145 lbs./acre or 0.25 lbs. AI/acre)
Transition (DE, KS, KY, MD, MO, VA, southern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY and RI)	2.45 lbs./1,000 sq. ft. (107 lbs./acre or 0.183 lbs. AI/acre)	3.33 lbs./1,000 sq. ft. (145 lbs./acre or 0.25 lbs. AI/acre)	4.9 lbs./1,000 sq. ft. (213 lbs./acre or 0.366 lbs. AI/acre)
South (AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN and TX)	3.33 lbs./1,000 sq. ft. (145 lbs./acre or 0.25 lbs. AI/acre)	4.9 lbs./1,000 sq. ft. (213 lbs./acre or 0.366 lbs. AI/acre)	6.67 lbs./1,000 sq. ft. (291 lbs./acre or 0.50 lbs. AI/acre)
Coastal South (HI, FL, southern coastal areas of AL, GA, LA, MS, NC, SC, and TX)	4.9 lbs./1,000 sq. ft. (213 lbs./acre or 0.366 lbs. AI/acre)	6.67 lbs./1,000 sq. ft. (291 lbs./acre or 0.50 lbs. AI/acre)	6.67 lbs./1,000 sq. ft.* (291 lbs./acre or 0.50 lbs. AI/acre)
* Split applications may be made 5-10 weeks apart.			

SPLIT OR SEQUENTIAL APPLICATIONS: Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with rates in the table above being split between two applications made 5-10 weeks apart. Where longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application provided the maximum application rate per year is not exceeded. Refer to the "Use Restrictions" section above.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS: SPRING APPLICATION: When used as directed for crabgrass control in spring, Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr will also control or suppress the following weeds when applied prior to their emergence:

GRASSES	
Common Name	Botanical Name
Barley, Barnyardgrass, Bluegrass (annual roughstalk), Brome, Crabgrass (large), Crabgrass (smooth), Crabgrass (Southern), Crowfootgrass, Dallisgrass (seedling), Foxtail (yellow & green), Goosegrass, Kikuyugrass*, Oats (wild), Ryegrass (annual & perennial), Sandbur, Smutgrass	Hordeum spp., Echinochloa crus-galli, Poa annua, Bromus spp., Digitaria sanguinalis, Digitaria ischaemum, Digitaria ciliaris, Dactyloctenium aegyptium, Paspalum dilatatum, Setaria faberi, Setaria Verdi, Eleusine indica, Pennisetum clandestinum, Avena fatua, Lolium spp., Cenchrus spp., Sporobolus indicus
BROADLEAF WEEDS	
Common Name	Botanical Name
Bittercress*, Carpetweed, Chickweed*, Geranium (Carolina*), Henbit, Knotweed (prostrate), Lespedeza (common)*, Marehail*, Medic (black), Mustard, Oxalis (buttercup), Parsley-piert*, Pigweed (redroot), Pineappleweed*, Purslane (common), Rocket (London), Shepherdspurse, Speedwell (common), Spurge (garden), Spurge (prostrate), Spurge (spotted), Woodsorrel (creeping), Woodsorrel (yellow)	Cardamine spp., Mullugo verticillata, Stellaria spp., Geranium carolinianum, Lamium spp., Polygonum aviculare, Lespedeza striata, Conyza canadensis, Medicago lupulina, Brassica spp., Oxalis pes-caprae, Alchemilla arvensis, Amaranthus retroflexus, Matricaria matricarioides, Portulaca oleracea, Sisymbrium irio, Capsella bursa-pastoris, Veronica arvensis, Euphorbia hirta, Euphorbia humistrata, Euphorbia maculata, Oxalis corniculata, Oxalis stricta
* Suppression only.	

FALL APPLICATION:When used as directed for late summer or early fall use, for crabgrass control through the early part of next spring, Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr will also control or suppress the following weeds when applied prior to their emergence:

WEEDS CONTROLLED	
Common Name	Botanical Name
Bluegrass (annual), Bittercress, Chickweed, Geranium (Carolina), Henbit, Parsley-piert, Pineappleweed, Shepherdspurse	Poa annua, Cardamine spp., Stellaria spp., Geranium carolinianum, Lamium spp., Alchemilla arvensis, Matricaria matricarioides, Capsella bursa-pastoris

APPLICATION EQUIPMENT AND INSTRUCTIONS:This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference.

Apply this product uniformly over the treatment area. More uniformity can be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Check equipment frequently to verify calibration and proper functioning. Clean and lubricate spreader after use. Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr provides preemergence control or suppression of annual grass and broadleaf weeds in landscape ornamentals. Refer to "For Use on Turf" section of this label for a complete listing of weeds controlled or suppressed. This product may be applied to soil

surfaces or mulch where ornamental plants are grown for aesthetic purposes in ornamental gardens, parks, golf courses, and residential areas.

APPLICATION INSTRUCTIONS: Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination to bare ground or mulch. For best results, apply to soil that is free of clods, weeds and debris such as leaves. Before application, any existing vegetation should be removed by hand weeding, cultivation, or with the use of a postemergence herbicide. Once the application is made, do not disturb the soil surface as the herbicide barrier will be broken. Irrigate immediately after application to remove herbicide particles from the foliage to prevent foliar damage to ornamentals.

SPLIT OR SEQUENTIAL APPLICATIONS: Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with rates in Table 2 being split between two applications made 5-10 weeks apart. Where longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application provided the maximum application rate per year is not exceeded.

- USE RESTRICTIONS
- Only apply this product to established ornamentals.
  - DO NOT apply this product directly to the bare roots of ornamental plants as injury may result.
  - DO NOT incorporate this product into soil. Dilution of active ingredient and plant root injury may occur.
  - DO NOT apply to soil around ornamentals that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes, or disease.
  - DO NOT apply under conditions that would affect product distribution (such as windy conditions).
  - DO NOT apply this product on grasses grown for seed.
  - DO NOT graze livestock or feed forage cut from areas treated with this product.

USE RATESApply Sure Green™ Grass & Weed Barrier™ Plus Lawn Food Containing Dithiopyr with a properly calibrated spreader that will provide accurate, even particle distribution. Refer to Table 2 below for product rates when treating a small area. DO NOT apply more than 6.67 lbs. of this product per 1,000 sq. ft., per application and no more than 20 lbs. of this product per 1,000 sq. ft. per year.

Table 2: Application Rates For Landscape Ornamentals		
LBS PRODUCT/ACRE	LBS PRODUCT/1,000 SQ. FT.	LBS PRODUCT/100 SQ. FT.
291	6.67	0.667 (10.73 oz.)

APPLICATION EQUIPMENT AND INSTRUCTIONS:This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference.

Apply this product uniformly over the treatment area. More uniformity can be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Check equipment frequently to verify calibration and proper functioning. Clean and lubricate spreader after use.

TOLERANT ORNAMENTALSWhen applied as directed under the conditions described on this label, the ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. A limited area involving only a few plants should be treated for tolerance evaluations before large-scale applications.

See other side for: Table 2. Tolerant Ornamental Species – All States Except CA

GUARANTEED ANALYSIS

18-3-3

Total Nitrogen (N) .....18.00%  
18.00% Ammoniacal Nitrogen

Available Phosphate (P2O5).....3.00%  
Soluble Potash (K2O) .....3.00%  
Sulfur (S) .....14.00%  
14.00% Combined Sulfur (S)

Iron (Fe)..... 1.00%

Derived from: Ammonium Sulfate, Ammonium Phosphate, Urea Muriate of Potash, and Iron Oxide.

Information concerning the raw materials composing this product can be obtained by writing to Spra-Green Lawn Fertilizer, LLC and referring to the product name.

8 51645 00704 2

Manufactured/Guaranteed by:  
**Spra-Green Lawn Fertilizer, LLC**  
PO BOX 845 • 221 N. Main • Portales, NM 88130



Common Name	Botanical Name	Tolerant Cultivars
Azalea	Rhododendron spp.	Brilliant, Buceanese, Carro, Chimes (Belgian), Elsie Lee, Exbury, Fashion, Heazler Beauty, Hershey Red, Higgas, Hinoconsmoon, Holland (Hybrid), Marion Lee, Northern Lights, Orange Cup, Orchid Lights, Snow, Southern Charm
Azalea, Flame	Rhododendron calendulaceum	
Azalea, Krishna	Nandina domestica	
Bamboo, Heavenly	Barberis thunbergii	Aurea, Dwarf Pygmy, Green, Kobold, Pygmy Red, Rose Glow
Basket Flower	Barberis vulgaris	
Beeberry (common)	Arctostaphylos uva-ursi	Massachusetts
Birch, River	Betula nigra	
Blackeyed Susan	Rudbeckia hirta	
Blanket Flower	Gallardia spp.	Goldstrum
Blueberry	Vaccinium spp.	Bluecrop, Blue Jay, Jersey, North Blue, Northland
Boxwood, Japanese	Buxus microphylla	Japanica
Boxwood, Weiler	Buxus sempervirens	Moonglight
Broom	Gentia plicata	Vancouver Gold
Bugle Carpet	Camellia japonica	Debutante, Mathotiana Supreme, Chansonette
Candy Tuft	Carex spp.	Snow White
Cedar, Red	Juniperus virginiana	
Celastrus	Celastrus spp.	
Cockscomb, Plumosa	Centaurea montana	Scarlet Plumosa
Coleus	Coleus blumei	Red Kewpie
Columbine	Aquilegia spp.	
Copper Leaf	Acalypha wilkesiana	Moonbloom
Coreopsis	Coreopsis spp.	
Cornflower	Centaurea spp.	
Cotoneaster	Cotoneaster apiculatus	
Coyotebush	Baccharis peltata	
Cycads	Cycads revolute	
Cypress, Bald	Taxodium distichum	
Cypress, Italian	Cupressus sempervirens	Glauca
Cypress, Japanese Falis	Chamaecyparis obtuse	Gracilis
Cupress, Leyland	Cupressocyparis leylandii	King Alfred
Dafnolil	Narcissus spp.	
Daylily	Hemerocallis spp.	Aztec Gold, Bright Yellow, (Hybrid), Single Gold (Evergreen), Wilson's Yellow
Delphinium	Delphinium spp.	Magic Fountain
Dianthus (Sweet William)	Cornus florida	
Dogwood, American	Cornus sericea	Flavirammas
Dusty Miller	Senecio cineraria	
Elm, Drake	Ulmus parvifolia	
Euonymus	Euonymus alatus	Argenteo-variegata, Aureo-marginata, Colorata, Emerald Gaiety, Emerald 'N Gold, Gold Edge, Gold Princess, Silver King, Tricolor, Vegetus
Fan Palm, European	Chamaeceros humilis	
Fern (various)	Asparagus spp.	
Fescue	Festuca glauca	Rainbow
Fetterbush	Leucothoe fontinalis	
Ficus	Ficus religiosa	Nitida
Fir, Fraser	Abies fraseri	
Forythia	Forsythia X intermedia	Arnold Dwarf, Bronxensis Dwarf, Lynnwood Gold, Meadowlark, Weeping
Fuchsia	Fuchsia spp.	
Gallum	Gallum odoratum	
Gardenia	Gardenia jasminoides	Mystery, Radicans
Geranium	Pelargonium X hortorum	
Gum	Eucalyptus citrodora	
Hawthorn	Crataegus spp.	Cockspur White, Crimson Cloud, Enchantress, Jack Evans, Washington White
Heather, Twisted	Erica cinerea	Mediterranean Pink
Hemlock, Canada	Tsuga canadensis	
Hibiscus	Hibiscus spp.	Blue Bird, Brilliant, Hula Girl
Holly	Ilex spp.	Blue Boy, Blue Girl
Ilex X meserveae	Ilex X meserveae	Burdott, China Girl, Compact, Forsteri, Helleri, Japanese Northern Beauty, Needpoint, Nellie R. Stevens
Ilex X attenuata	Ilex X attenuata	Savannah

Common Name	Botanical Name	Tolerant Cultivars
Snapdragon	<i>Antirrhinum</i> spp.	Dragon Blood Red, Red Carpet, Yellow
Sourwood	<i>Oxydendrum arborescens</i>	
Spiraea	<i>Aristida x arandensis</i>	
Spiraea	<i>Spiraea</i> spp.	Anthony Water Red, Dechthica, Goldenflame Red, Snowmound White.
Spiraea, Garland	<i>Spiraea</i> spp.	
Spiraea, Black Hills	<i>Picea glauca</i>	
Spruce, Colorado Blue	<i>Picea pungens</i>	
Spruce, Norway	<i>Picea abies</i>	
Sweetflag, Grassly-Le	<i>Picea glauca</i>	Conica
Sweetgum	<i>Liquidambar styraciflua</i>	
Sycamore	<i>Platanus occidentalis</i>	
Tree Fern (Tiki Fern)	<i>Asparagus virgatus</i>	
Trumpet Flower, Evening	<i>Ipomoea pes-caprae</i>	
Tulip	<i>Tulipa</i> spp.	Apeldoorn
Verbena, Shrub	<i>Lantana setowiana</i>	St. Paul
Vervain	<i>Veronica</i> spp.	
Viburnum	<i>Viburnum</i> spp.	American Cranberry Bush, Arrowwood, Common Snowball, European Cranberry Bush, Linden, Mohican, Wright
Vinca (Periwinkle)	<i>Vinca minor</i>	
Windmill Palm	<i>Trachycarpus fortunei</i>	
Xylosma	<i>Xylosma congestum</i>	
Yarrow	<i>Achillea</i> spp.	Dwarf
Yaupon	<i>Ilex vomitoria</i>	
Yew	<i>Taxus cuspidata</i>	Densitormis
	<i>Taxus x media</i>	
* Ornamental species only. Do not use on food producing trees and ornamentals.		

STORAGE AND DISPOSAL		
Do not contaminate water, food or feed by storage or disposal.		
STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Empty, non-refillable containers, do not reuse or refill this container. Offer for recycling, if available. Do not reuse unless required for recycling. If recycling is not available, then dispose of container in a sanitary landfill or, if allowed by State and local authorities by burning, if burned, stay out of smoke. If partly filled: Call your local solid waste agency for disposal instructions. Never pour unused product down any indoor or outdoor drain.		
CONDITIONS OF SALE AND WARRANTY		
NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying and using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.		
The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Lawn and turf injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Spira-Green Lawn Fertilizer, LLC or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.		
Spira-Green Lawn Fertilizer, LLC warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.		
TO THE EXTENT PERMITTED BY LAW, SPIRA-GREEN LAWN FERTILIZER, LLC, NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL, DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF, THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.		
and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.		